

Application Serial: 09/813,197

**VERSION WITH MARKINGS TO SHOW CHANGES MADE**  
(Response to File the Notice of Missing Parts)

The paragraph beginning at line 22 on page 19 has been amended as follows:

- - Figure 11[(A)] provides examples of non-native amino acids with reporter properties, [Figure 11(B) shows] illustrates participation of a reporter in protein synthesis, and [Figure 11(C) shows] illustrates synthesis of a reporter.- -

The paragraph beginning at line 14 on page 51 has been amended as follows:

- - The chemical synthesis of a reporter can be based on the linkage of a chemical moiety or a molecular component having reporter properties with a native amino acid residue. There are many fluorescent molecules which are sensitive to their environment and undergo a change in the wavelength of emitted light and yield of fluorescence. When these chemical moieties, coupled to amino acids, are incorporated into the synthesized protein, their environments are altered because of a difference between the bulk aqueous medium and the interior of a protein which can causes reduced accessibility to water, exposure to charged ionic groups, reduced mobility, and altered dielectric constant of the surrounding medium. Two such examples are shown in Figure 11[A].- -

The paragraph beginning at line 14 on page 52, has been amended as follows:

A second example of a reporter is a marker based on coumarin such as 6,7-(4', 5'-proline)coumarin. This compound can be chemically synthesized by coupling a fluorophore like coumarin with an amino-acid structural element in such a way that the fluorophore would alter its emission or absorption properties after forming a peptide linkage (Figure 11[B]). For example, a proline ring containing secondary amino functions will participate in peptide bond formation similar to a normal primary amino group. Changes in fluorescence occur due to the co-planarity of the newly formed peptide group in relation to the existing fluorophore. This increases conjugation/delocalization due to the  $\pi$ -electrons of nitrogen-lone pair and carbonyl-

***PATENT***

Attorney Docket No. AMBER-06311

group in the peptide bond. Synthesis of such compounds is based on coumarin synthesis using ethylacetoacetate (Figure 11[C]).